

## **Magnetic Resonance Imaging (MRI) – Prostate**

### **What is Magnetic Resonance Imaging (MRI) of the Prostate?**

#### **Definition**

Magnetic Resonance Imaging (an MRI) of the prostate gland in men creates images of this gland and its surrounding structures in order to evaluate their condition and to assist with the diagnosis and treatment of problems. In some ways, an MRI is similar to an x-ray, but it is also different in several important ways. Ultrasound and MRI are the most common techniques used to create images of the prostate.

#### **How It Works**

During an MRI procedure on the body, a contrast material (dye) may be injected into the body via an IV (intravenous drip). The contrast material injected through the IV contains a magnetic substance, which gets distributed throughout the body via the blood stream. When the MRI equipment is put in motion, the contrast material reacts to the magnets to reveal the details of the structures within and around the prostate, similar to the way x-rays create images of bones. The difference is an MRI uses a powerful magnetic field and radiofrequency pulses to create detailed pictures, whereas an x-ray uses radiation. X-rays are usually less effective for creating images of soft organs and tissues in the body because this soft matter shows up as shades of grey, unlike bones which are bright white and can be seen in detail. In most cases, an MRI can also produce more detailed images of the prostate than other methods.

#### **Common Uses**

MRIs of the prostate are commonly used to:

- Diagnose infection, an enlarged prostate, or abscesses
- Detect and evaluate prostate cancer
- Determine if prostate cancer has spread beyond the walls of the prostate gland into other parts of the body

#### **Benefits and Risks**

##### ***Benefits***

- Magnetic resonance imaging (MRIs) can help identify causes of pain and help the patient make informed decisions about further treatment.
- Magnetic resonance imaging (MRIs) can produce much higher quality images of the prostate than other methods, which can make a crucial difference in a doctor's ability to quickly and accurately evaluate the cause of the patient's discomfort.
- Magnetic resonance imaging is a non-invasive procedure.
- The patient is not exposed to any x-ray radiation during an MRI procedure.
- The contrast material (dye) used in MRIs is less likely to cause allergic reactions than those used for x-rays and CT scans.

##### ***Risks***

- There is an extremely small risk of developing an infection (less than 1 in 1,000) at the site where the patient's skin is punctured to establish the intravenous drip (IV), which would require treatment with antibiotics if it were to arise.
- Although rare, few patients may experience side effects from the contrast material. Patients are instructed to notify the technician immediately if any side effects are noted during the procedure.
- Medical devices inside the body may cause problems during any MRI exam because they will be affected by the magnets in the MRI equipment. Therefore, patients are carefully screened to insure it is safe to have an MRI scan.
- In extremely rare cases, patients with compromised kidney function who are injected with high doses of gadolinium contrast material (magnetic dye) during an MRI can develop nephrogenic systemic fibrosis.
- Other potential risks may vary from patient to patient; the patient should speak to his or doctor before the procedure about any questions or concerns.

### **How Should I Prepare for My Appointment?**

Restrictions on what a patient may eat or drink before an MRI procedure vary, based on the type of procedure to be performed, the specific body parts being and the facility's guidelines. In certain cases, the patient may be asked to fast (to avoid consuming any food or liquids aside from water) for 8-12 hours before the procedure. The patient should check with the facility to see if he/she can eat, drink, and take medication as usual before the procedure. The patient should notify his doctor of any drugs or materials to which he is allergic and should notify doctor of any other pertinent details of his medical history (prescriptions, recent illnesses or injuries, or serious health problems, etc.).

### **What Will Happen During the Procedure?**

- The patient will be asked to remove any items of clothing, jewelry, or other accessories that might interfere with the procedure.
- A patient gown will be provided unless the patient is wearing clothing that is free of metal (zippers, buttons, snaps, decorative glitter, etc.).
- Because the scans are loud, hearing protection will be provided.
- If contrast has been ordered, the technician may instruct the patient to lie down on the examination table and slide the table inside the cylindrical chamber to take some initial images of the prostate before the contrast material is introduced into the body's system (i.e., taking "before" photos).
- The technician will insert a needle into a vein to establish an IV (a harmless saline solution would most likely be dripping from the IV bag until it is time to insert the contrast dye via the IV, after any initial scans).
- The patient will be positioned on the moveable examination table. Then, the examination table will be slid into position inside the cylindrical tube.
- The patient will need to remain very still while the MRI machine takes images of the prostate.
- The patient will be alone in the exam room (a parent or friend may be allowed to stay in the room), but the technician will be able to see and speak with the patient through a two-way intercom. The MRI machine will be noisy (it "buzzes" and "hums" while the magnets do the scanning).

- After all the necessary images have been collected (sometimes several “runs” are necessary to obtain enough images of the body part or area in question), the exam table will be moved back out of the cylindrical tube and the patient may get up.
- If an IV was inserted, it would be removed after the procedure is complete.

**What Should I Expect After the Procedure?**

If the patient has been sedated as part of the MRI procedure, a short recovery period may be required; otherwise, the patient may generally go on with his normal routine and activities.